


## Benzotrichloride

### CAS No. 98-07-7

Reasonably anticipated to be a human carcinogen  
First Listed in the *Fourth Annual Report on Carcinogens* (1985)



### Carcinogenicity

Benzotrichloride is *reasonably anticipated to be a human carcinogen* based on sufficient evidence of carcinogenicity in experimental animals (IARC 1982a,b, 1987, 1999). When administered by gavage, benzotrichloride induced squamous cell carcinomas of the forestomach and adenocarcinomas of the lung in female mice. Benzotrichloride was tested in three studies by skin application to female mice. It induced squamous cell carcinomas of the skin and lung in all three experiments, and upper digestive tract tumors (carcinomas of the lips, tongue, esophagus, or stomach) and lymphomas were observed in two of the three experiments. In a mouse-lung tumor bioassay, benzotrichloride increased the incidence of lung adenomas.

No data were available to evaluate the carcinogenicity of  $\alpha$ -chlorinated toluenes, including benzotrichloride, in humans (IARC 1982a,b, 1987, 1999). There is evidence that employment in the production of chlorinated toluenes, which involves potential exposure to benzotrichloride as well as other chemicals, increases the risk of respiratory cancer.

### Properties

Benzotrichloride is a clear, colorless to yellowish, oily liquid with a penetrating odor. It is insoluble in water and soluble in benzene, diethyl ether, and ethanol. Benzotrichloride is unstable and hydrolyzes to benzoic acid and hydrochloric acid in the presence of moisture (IARC 1999). It is available in the United States as a liquid containing 99% active ingredient.

### Use

Benzotrichloride is used extensively as a chemical intermediate. Its most important derivatives are benzoyl chloride and substituted benzophenones used to stabilize plastics in the presence of ultraviolet light (SRI 1986, IARC 1999). It is also used as a dye intermediate in the preparation of eight dyes and pigments, including five produced in commercial quantities in the United States. Additionally, benzotrichloride is used to make benzotrifluoride, hydroxybenzophenone, antiseptics, and antimicrobial agents (IARC 1982a).

### Production

The 1997 *Directory of Chemical Producers* listed one producer of benzotrichloride (SRI 1997). Chem Sources (2001) identified eight current U.S. suppliers. From 1982 to 1990 and in 1992, the USITC identified two companies producing an unspecified amount of benzotrichloride (USITC 1983-1991, 1994). In 1991 and 1994, only one manufacturer was reported (USITC 1993, 1995). Currently, two significant U.S. producers of benzotrichloride with a capacity totaling 68 million lb were reported (Seper 2001). The latest production volume found was for western countries in 1988 when 69 million lb of the chemical were produced (IARC 1999). U.S. production volume for 1982 was 35 million lb (HSDB 2000).

In 1986, U.S. imports of benzotrichloride totaled 363,000 lb (HSDB 2000). In 1983, imports through the principal customs districts totaled 39,600 lb (USITC 1984b). The 1979 TSCA Inventory reported three companies with a total production of 30 million lb in

1977 (TSCA 1979). It was estimated that approximately 40 million lb of benzotrichloride are needed annually for the U.S. production of benzoyl chloride alone. Benzotrichloride has been produced commercially in the United States since at least 1919 (IARC 1982a).

### Exposure

The primary routes of potential human exposure to benzotrichloride are inhalation, ingestion, and dermal contact. Its exclusive use as a chemical intermediate results in minimal potential consumer exposure. Exposure to the chemical from industrial fugitive emissions is also expected to be very low due to its ability to hydrolyze rapidly in the presence of moisture. Consumer exposure to small amounts may occur during the use of pharmaceuticals made with benzotrichloride. Potential occupational exposure of workers could be significant due to releases in the work environment in the liquid or vapor form. A realistic exposure assessment is not possible because the number of workers, exposure levels, and releases to the environment are not documented (CHIP 1982). The National Occupational Exposure Survey (1981-1983) estimated that 171 workers potentially were exposed to benzotrichloride (NIOSH 1984). Potential occupational exposure can occur during the production, formulation, packaging, and application of antiseptics made with benzotrichloride or benzoyl chloride. Benzotrichloride has been identified in surface waters at unreported concentrations. It does not occur naturally (IARC 1982a). EPA's Toxic Chemical Release Inventory (TRI) estimated that 2,175 lb of benzotrichloride were released to the air by seven facilities that produced, processed, or used the chemical in the United States in 1999 (TRI99 2001).

### Regulations

#### DOT

Benzotrichloride is considered a hazardous material and special requirements have been set for marking, labeling, and transporting this material

#### EPA

##### Clean Air Act

NESHAP: Listed as a Hazardous Air Pollutant (HAP)

NSPS: Manufacture of substance is subject to certain provisions for the control of Volatile Organic Compound (VOC) emissions

##### Comprehensive Environmental Response, Compensation, and Liability Act

Reportable Quantity (RQ) = 10 lb

##### Emergency Planning and Community Right-to-Know Act

Toxics Release Inventory: Listed substance subject to reporting requirements

Threshold Planning Quantity (TPQ) = 100 lb

Reportable Quantity (RQ) = 10 lb

##### Resource Conservation and Recovery Act

Listed Hazardous Waste: Waste codes in which listing is based wholly or partly on substance - U023, K015, K149

Listed as a Hazardous Constituent of Waste

### Guidelines

#### ACGIH

Threshold Limit Value - Ceiling (TLV-C) = 0.1 ppm

### REFERENCES

- ChemSources. 2001. Chemical Sources International, Inc. <http://www.chemsources.com>.
- CHIP. 1982. Chemical Hazard Information Profile. Benzotrichloride. Washington, D.C.: U.S. Environmental Protection Agency, Office of Pesticide Programs and Toxic Substances.
- HSDB. 2000. Hazardous Substances Data Base. National Library of Medicine. <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>.
- IARC. 1982b. Chemicals, Industrial Processes and Industries Associated with Cancer in Humans. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, Supplement 4. Lyon, France: International Agency for Research on Cancer. 292 pp.
- IARC. 1982a. Some Industrial Chemicals and Dyestuffs. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 29. Lyon, France: International Agency for Research on Cancer. 416 pp.
- IARC. 1987. Overall Evaluations of Carcinogenicity. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, Supplement 7. Lyon, France: International Agency for Research on Cancer. 440 pp.
- IARC. 1999. Re-evaluation of Some Organic Chemicals, Hydrazine, and Hydrogen Peroxide. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans, vol. 71. Lyon, France: International Agency for Research on Cancer. 1589 pp.
- NIOSH. 1984. National Occupational Exposure Survey (1981-83). Cincinnati, OH: U. S. Department of Health and Human Services. <http://www.cdc.gov/noes/noes3/empl0003.html>.
- Seper, K. 2001. Benzyl Chloride, Benzal Chloride and Benzotrichloride. In Kirk-Othmer Encyclopedia of

## *SUBSTANCE PROFILES*

- Chemical Technology. New York: John Wiley and Sons.
- SRI. 1986. Directory of Chemical Producers, United States, 1985. Stanford Research Institute, Menlo Park, CA: SRI International.
- SRI. 1997. Directory of Chemical Producers, United States, 1997. Stanford Research Institute, Menlo Park, CA: SRI International.
- TRI99. 2001. Toxic Chemical Release Inventory 1999. Data contained in the Toxic Chemical Release Inventory (TRI). National Library of Medicine. <http://www.epa.gov/triexplorer/>.
- TSCA. 1979. Toxic Substances Control Act, Chemical Substances Inventory.
- USITC. 1983. Synthetic Organic Chemicals, United States Production and Sales, 1982. USITC Publication No 1422. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1984. Imports of Benzenoid Chemicals and Products, 1983. USITC Publication No 1548. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1984. Synthetic Organic Chemicals, United States Production and Sales, 1983. USITC Publication No 1588. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1985. Synthetic Organic Chemicals, United States Production and Sales, 1984. USITC Publication No 1745. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1986. Synthetic Organic Chemicals, United States Production and Sales, 1985. USITC Publication No 1892. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1987. Synthetic Organic Chemicals, United States Production and Sales, 1986. USITC Publication No 2009. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1988. Synthetic Organic Chemicals, United States Production and Sales, 1987. USITC Publication No 2118. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1989. Synthetic Organic Chemicals, United States Production and Sales, 1988. USITC Publication No 2219. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1990. Synthetic Organic Chemicals, United States Production and Sales, 1989. USITC Publication No 2338. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1991. Synthetic Organic Chemicals, United States Production and Sales, 1990. USITC Publication No 2470. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1993. Synthetic Organic Chemicals, United States Production and Sales, 1991. USITC Publication No 2607. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1994. Synthetic Organic Chemicals, United States Production and Sales, 1992. USITC Publication No 2720. Washington, D.C.: U.S. Government Printing Office.
- USITC. 1995. Synthetic Organic Chemicals, United States Production and Sales, 1994. USITC Publication No 2933. Washington, D.C.: U.S. Government Printing Office.